# **Medicare Advantage Medical Policy**



An independent licensee of the Blue Cross and Blue Shield Association

> \*Current Policy Effective Date: 11/1/22 (See policy history boxes for previous effective dates)

# Title: Spinal Surgery: Automated Percutaneous and Endoscopic Discectomy

## **Description/Background**

Traditionally, discectomy and microdiscectomy are performed manually through an open incision. Percutaneous discectomy describes techniques by which disc decompression is accomplished by the physical removal of disc material rather than its ablation. These techniques have been modified by the use of automated devices that involve placement of a probe within the intervertebral disc and aspiration of disc material using a suction cutting device. Removal of disc herniations under endoscopic visualization is also being investigated.

Back pain or radiculopathy related to herniated discs is an extremely common condition and a frequent cause of chronic disability. Although many cases of acute low back pain and radiculopathy will resolve with conservative care, a surgical decompression is often considered when the pain is unimproved after several months and is clearly neuropathic in origin, resulting from irritation of the nerve roots. Open surgical treatment typically consists of discectomy in which the extruding disc material is excised. When performed with an operating microscope, the procedure is known as microdiscectomy.

Minimally invasive options have also been researched, in which some portion of the disc material is removed or ablated, although these techniques are not precisely targeted at the offending extruding disc material. Ablative techniques include laser discectomy and radiofrequency (RF) decompression. In addition, intradiscal electrothermal annuloplasty is another minimally invasive approach to low back pain. In this technique, RF energy is used to treat the surrounding disc annulus.

This policy addresses automated percutaneous and endoscopic discectomy, in which the disc decompression is accomplished by the physical removal of disc material rather than its ablation. A discectomy is performed manually through an open incision, using cutting forceps to remove

nuclear material from within the disc annulus. This technique has been modified by automated devices that involve placement of a probe within the intervertebral disc and aspiration of disc material using a suction cutting device. Endoscopic techniques may be intradiscal or may involve the extraction of non-contained and sequestered disc fragments from inside the spinal canal using an interlaminar or transforaminal approach. Following insertion of the endoscope, the decompression is performed under visual control.

### **Regulatory Status:**

The Stryker DeKompressor® Percutaneous Discectomy Probe (Stryker) and the Nucleotome® (Clarus Medical) are examples of percutaneous discectomy devices that received clearance from the U.S. Food and Drug Administration (FDA) through the 510(k) process. Both have the same labeled intended use, i.e., "for use in aspiration of disc material during percutaneous discectomies in the lumbar, thoracic and cervical regions of the spine." FDA product code: HRX.

A variety of endoscopes and associated surgical instruments have received marketing clearance through the FDA's 510(k) process.

## **Medical Policy Statement**

Automated percutaneous discectomy and endoscopic discectomy are considered experimental/investigational. There is insufficient evidence obtained from well-designed and executed randomized controlled trials to evaluate the impact of automated percutaneous discectomy and endoscopic discectomy on net health outcome.

# **Inclusionary and Exclusionary Guidelines** (Clinically based guidelines that may support individual consideration and pre-authorization decisions)

N/A

**CPT/HCPCS Level II Codes** (Note: The inclusion of a code in this list is not a guarantee of coverage. Please refer to the medical policy statement to determine the status of a given procedure)

Established codes:

N/A

Other codes (investigational, not medically necessary, etc.):

62287 62380 C9757

## Rationale

Evidence reviews assess the clinical evidence to determine whether the use of a technology improves the net health outcome. Broadly defined, health outcomes are length of life, quality of life, and ability to function-including benefits and harms. Every clinical condition has specific outcomes that are important to patients and to managing the course of that condition. Validated outcome measures are necessary to ascertain whether a condition improves or worsens; and whether the magnitude of that change is clinically significant. The net health outcome is a balance of benefits and harms.

To assess whether the evidence is sufficient to draw conclusions about the net health outcome of a technology, 2 domains are examined: the relevance and the quality and credibility. To be relevant, studies must represent one or more intended clinical use of the technology in the intended population and compare an effective and appropriate alternative at a comparable intensity. For some conditions, the alternative will be supportive care or surveillance. The quality and credibility of the evidence depend on study design and conduct, minimizing bias and confounding that can generate incorrect findings. The randomized controlled trial (RCT) is preferred to assess efficacy; however, in some circumstances, nonrandomized studies may be adequate. RCTs are rarely large enough or long enough to capture less common adverse events and long-term effects. Other types of studies can be used for these purposes and to assess generalizability to broader clinical populations and settings of clinical practice.

### AUTOMATED PERCUTANEOUS DISCECTOMY

#### **Clinical Context and Therapy Purpose**

The purpose of automated percutaneous discectomy in individuals who have herniated intervertebral disc(s) is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does the use of automated percutaneous discectomy improve the net health outcome in individuals with herniated intervertebral disc(s)?

The following **PICOs** were used to select literature to inform this review.

#### **Populations**

The relevant population of interest is individuals with herniated intervertebral disc(s).

#### Interventions

The therapy being considered is automated percutaneous discectomy.

Percutaneous discectomy is provided in a hospital setting with specialized staff and equipped to perform the surgical procedure and post-surgical care.

#### Comparators

The following therapies and practices are currently being used to treat herniated intervertebral disc(s): conservative therapy and open discectomy or microdiscectomy.

### Outcomes

The general outcomes of interest are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Specific outcomes measured by specific instruments include improvements in functional outcomes assessed on the Oswestry Disability Index (ODI), reductions in pain using a visual analog scale (VAS), improvements in quality of life measured on the 36-Item Short-Form Health <u>Survey</u> (SF-36) and Euro-QOL-5D, and reductions in medication usage.

To assess outcomes, follow-up at 1 year is considered appropriate.

#### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.

### **Systematic Reviews**

A 2015 systematic review and network meta-analysis by Lewis et al compared trials of 21 different treatment strategies for sciatica.<sup>1</sup> Examples of the 21 treatment strategies included in the analysis are conservative care, disc surgery, intraoperative interventions, epidural injections, biologic agents, and percutaneous discectomy. Under the category of "percutaneous discectomy," reviewers combined automated percutaneous discectomy, percutaneous automated nucleotomy, nucleoplasty, and laser discectomy. They searched 28 databases and trial registries through December 2009. Ninety studies were included and 10 involved the percutaneous discectomy category as an intervention. Of the 10, 4 are relevant to this evidence review: 2 case-control studies of percutaneous endoscopic discectomy (2006, 2007), 1 RCT of percutaneous endoscopic discectomy (1993), and 1 RCT of automated percutaneous discectomy (1995). The remaining studies were published in a foreign language or involved the comparators nucleolysis and chemonucleolysis, rather than open discectomy or microdiscectomy. The global effects odds ratio for the category of percutaneous discectomy compared with inactive control was 0.82 (95% confidence interval [CI], 0.39 to 1.72), which was inferior to disc surgery, epidural injections, and intraoperative interventions. The pain intensity weighted mean difference for the category of percutaneous discectomy compared with inactive control was 11.5 (95% CI, -18.6 to 41.6). Reviewers concluded that there was no support for the effectiveness of percutaneous discectomy for the treatment of sciatica. Due to the inclusion of additional interventions into the broad category of percutaneous discectomy in this review, the relevance of these results for our evidence review is limited.

#### **Randomized Controlled Trials**

The 2002 LAPDOG trial compared automated percutaneous discectomy with open discectomy in patients with lumbar disc herniation.<sup>2</sup> No additional RCTs have been identified since the 2002 LAPDOG trial. The trial was designed to recruit 330 patients but enrolled 36 patients for reasons not readily apparent. Twenty-seven patients were available at follow-up, with efficacy reported by 41% of those undergoing automated percutaneous discectomy and by 40% of those undergoing conventional discectomy. The trialists concluded that "It is difficult to understand the remarkable persistence of percutaneous discectomy in the face of a virtually

complete lack of scientific support for its effectiveness in treated lumbar disc herniation." The tables below more fully describe key characteristics, results, and limitations of the LAPDOG trial.

Table 1. Characteristics	s of the LAPDOG Trial
--------------------------	-----------------------

Study Cou	untries	Sites	Dates	Participants	Interventions
Haines et US. al (2002) <sup>2</sup> Car	., nada	10	NR	Patients with predominantly unilateral leg pain or paresthesia with no previous treatment for lumbar spinal disease, at least 2 of 4 objective signs, and an imaging study confirming disc herniation at the appropriate level	Automated percutaneous discectomy vs. conventional discectomy

LAPDOG: Lumbar Automated Percutaneous Discectomy Outcomes Group; NR: not reported.

#### Table 2. Results of the LAPDOG Trial

Success <sup>a</sup> (at	Treatment Failure <sup>b</sup> (at	SF-36 Physical Functioning	SF-36 General Health Subscore	Modified Roland Score
6 months)	6 months)	Subscore		
2				
27	27	NR	NR	NR
7 (41%)	10 (59%)	Pre- vs. postoperative mean difference: 35.7	Pre- vs. postoperative mean difference: 5.0	Pre- vs. postoperative mean difference: 9.7
4 (40%)	6 (60%)	Pre- vs. postoperative mean difference: 36.1	Pre- vs. postoperative mean difference: 8.0	Pre- vs. postoperative mean difference: 10.6
0.95	0.95	0.96	0.58	0.74
2	Success <sup>a</sup> (at     6 months)     27     7 (41%)     4 (40%)     0.95	Fourier   Frequencies     Success <sup>a</sup> (at 6 months)   Failure <sup>b</sup> (at 6 months)     27   27     7 (41%)   10 (59%)     4 (40%)   6 (60%)     0.95   0.95	FourierFrequenciesFourtherSuccessa (at 6 months)Failureb (at 6 months)Functioning Subscore2727NR7 (41%)10 (59%)Pre- vs. postoperative mean difference: 35.74 (40%)6 (60%)Pre- vs. postoperative mean difference: 36.10.950.950.96	Fourier Buccessa (at 6 months)Failureb (at 6 months)Functioning SubscoreSF-36 General Health Subscore2727NRNR2727NRNR7 (41%)10 (59%)Pre- vs. postoperative mean difference: 35.7Pre- vs. postoperative mean difference: 5.0Pre- vs. postoperative mean difference: 36.14 (40%)6 (60%)Pre- vs. postoperative mean difference: 36.1Pre- vs. postoperative mean difference: 8.00.950.950.960.58

LAPDOG: Lumbar Automated Percutaneous Discectomy Outcomes Group; NR: not reported; SF-36: 36-Item Short-Form Health Survey. <sup>a</sup>Success was defined as either an excellent or good result as defined by an outcome matrix.

<sup>b</sup>Failure was defined as not achieving success or requiring a second procedure during the follow-up period.

#### Table 3. Study Relevance Limitations of the LAPDOG Trial

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Duration of Follow-Up <sup>e</sup>
Haines et al (2002)2	4. Investigators believed that study inclusion criteria reflected an existing population with lumbar disc disease; however, results from only 27 patients were eventually analyzed from a planned enrollment of 330 patients			4. Primary outcomes of "success" or "failure" largely subjective in nature; investigators admit that the outcome measurement tool used can not be precisely reproduced	1,2. Outcomes reported only for 6 months of follow-up; 12 month follow-up was achieved for only 19 patients and the study did not report any of these results

LAPDOG: Lumbar Automated Percutaneous Discectomy Outcomes Group.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup>Population key: 1. Intended use population unclear; 2. Clinical context is unclear; 3. Study population is unclear; 4. Study population not representative of intended use.

<sup>b</sup>Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4.Not the intervention of interest.

<sup>c</sup>Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively. <sup>d</sup>Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

<sup>e</sup>Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

#### Table 4. Study Design and Conduct Limitations of the LAPDOG Trial

Study	Allocation <sup>a</sup>	Blinding <sup>b</sup>	Selective Reporting <sup>c</sup>	Data Completeness⁴	Power <sup>e</sup>	Statistical <sup>f</sup>
Haines et al (2002)2		1,2. Blinding did not appear to occur		1. Of 34 initially randomized patients, 9 were lost to follow-up, 6 month follow- up data was obtained on only 27 patients, and 12 month follow- up data was obtained for only 19 patients	3. Power estimates led the investigators to plan enrollment of 330 patients in order to reliably identify a difference in success rate of 15% or greater; results were analyzed on 27 patients	1. Beyond the cursory discussion of lack of power, a discussion of the statistical analyses is nonexistent

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. <sup>a</sup>Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control

"Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup>Blinding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician. <sup>c</sup>Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.

<sup>d</sup>Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials). <sup>e</sup>Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important

difference. Statistical kay: 1. Analysis is not appropriate for outcome type: (a) continuous: (b) binary: (c) time to event: 2. Analysis is not appropriate

<sup>1</sup>Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

All published trials have focused on lumbar disc herniation. There were no RCTs of automated percutaneous discectomy for cervical or thoracic disc herniation. A review of the evidence from American Society of Interventional Pain Physicians (2013) noted that "even though Dekompressor [disc removal system] may be considered a new interventional modality, the early studies were published approximately 8 years ago. Consequently, one would expect that the technique's continued use would be supported by more recent, high-quality evaluations."<sup>3</sup>

#### Section Summary: Automated Percutaneous Discectomy

The evidence for automated percutaneous discectomy in individuals who have herniated intervertebral disc(s) includes small RCTs and systematic reviews. Evidence from small RCTs does not support the use of this procedure. Well-designed and executed RCTs are needed to determine the benefits and risks of this procedure.

#### PERCUTANEOUS ENDOSCOPIC DISCECTOMY

#### **Clinical Context and Therapy Purpose**

The purpose of percutaneous endoscopic discectomy in individuals who have herniated intervertebral disc(s) is to provide a treatment option that is an alternative to or an improvement on existing therapies.

The question addressed in this evidence review is: Does the use of percutaneous endoscopic discectomy improve the net health outcome in individuals with herniated intervertebral disc(s)?

The following **PICOs** were used to select literature to inform this review.

#### Populations

The relevant population of interest is individuals with herniated intervertebral disc(s).

#### Interventions

The therapy being considered is percutaneous endoscopic discectomy.

#### Comparators

The following therapies and practices are currently being used to treat herniated intervertebral disc(s): conservative therapy and open discectomy or microdiscectomy.

#### Outcomes

The general outcomes of interest are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Specific outcomes measured by specific instruments include improvements in functional outcomes assessed on the ODI, reductions in pain using a VAS, improvements in quality of life measured on the SF-36 and Euro-QOL-5D, and reductions in medication usage.

To assess outcomes, follow-up at 1 year is considered appropriate.

#### **Study Selection Criteria**

Methodologically credible studies were selected using the following principles:

- To assess efficacy outcomes, comparative controlled prospective trials were sought, with a preference for RCTs;
- In the absence of such trials, comparative observational studies were sought, with a preference for prospective studies.
- To assess longer term outcomes and adverse events, single-arm studies that capture longer periods of follow-up and/or larger populations were sought.
- Studies with duplicative or overlapping populations were excluded.

#### **Systematic Reviews**

A number of systematic reviews have evaluated the efficacy and safety of percutaneous endoscopic discectomy compared to open discectomy or microendoscopic discectomy. A comparison of the trials included in more recent systematic reviews (2016 to present) is shown in Table 5. The systematic reviews included a total of 67 trials published between 1993 and 2020. Characteristics and results of these reviews are summarized in Tables 6 and 7.

# Table 5. Trials Included in Systematic Reviews of Percutaneous Endoscopic Discectomy vs. Other Discectomy Procedures

Trials	Systematic Reviews	1							
	Cong et al (2016)4	Lietal (2016).5	Phan et al (2017)	Shi et al (2019)	Yu et al (2019) 8	Zhou et al (2020)9	Xu et al (2020) 10	Bai et al (2021)11	Gadjradj et al
	Cong et al (2010)	2010)	Than crui (2017)		ru et ul (2010)	2100 61 01 (2020)	Xu et ul (2020)	Dur et ul (2021)	(2021) <sup>12</sup>
Akcakaya et al (2016) <sup>13</sup>									•
Choi et al (2018) <sup>14.</sup>									•
Dai et al (2020) <sup>15.</sup>									•
Krappel et al (2001) <sup>16,</sup>									•
Tacconi et al (2019) <sup>17.</sup>									•
Tacconi et al (2020) <sup>18</sup> .									•
Tao et al (2018) <u>19.</u>									•
Wang et al (2017) <sup>20.</sup>									•
Xu et al (2020) <sup>21</sup> .									•
Ahn et al (2016) <sup>22.</sup>								•	
Chang et al (2018) <sup>23.</sup>								•	•
Liu et al (2017) <sup>24.</sup>								•	
Pan et al (2016) <sup>25.</sup>								•	•
Yao et al (2017) <sup>26.</sup>								•	
Yao et al (2017) <sup>27.</sup>								•	
Gibson et al (2017) <sup>28</sup>						•			•
Hsu et al (2013) <sup>29.</sup>						•			
Kim et al (2007) <sup>30.</sup>						•		•	
Qu et al (2017) <sup><u>31.</u></sup>						•			
Wang et al (2013) <sup>32.</sup>						•			
Zhao et al (2012) <sup>33.</sup>						•			
Yoon et al (2012) <sup>34.</sup>			•	•	•		•		
Li et al (2015) <sup>35.</sup>			•				•		
Sinkemani et al (2015) <sup>36.</sup>			•	•	•		•		
Song et al (2017) <sup>37,</sup>				•	•		•		
Tu et al (2017) <sup>38.</sup>							•		
Liu et al (2018) <sup>39,</sup>				•	•	•	•	•	
Li et al (2018) <u>40.</u>				•	•	•	•		
Abdurexiti et al (2018) <sup>41</sup> .				•	•		•		
Chen et al (2018) <sup>42.</sup>				•	•	•	•	•	
Liu et al (2012) <sup><u>43</u>.</sup>					•				
Wu et al (2009)44.				•					
Yang et al (2015) <sup>45.</sup>				•		•			
Duan et al (2016) <sup>46</sup> .				•					
Zhao et al (2016)47.				•					
Ding et al (2017)48.				•					
Li et al (2017) <sup>49</sup> .				•					
Liu et al (2017) <sup>50,</sup>				•					
Luo et al (2017) <u>51.</u>				•					
Qu et al (2017) <sup>52.</sup>				•					
Chen et al (2018) <sup>53.</sup>				•					
Wu et al (2018) <sup>54.</sup>				•					
Belykh et al (2016)55.				•					
Chen et al (2015) <sup>56.</sup>			•						
Choi et al (2016)57.			•			•			
Garg et al (2011) <sup>58</sup> .	•		•						
Hermantin et al (1999)59.	•		•						٠
Huang et al (2005) <sup>60.</sup>			•						
Hussein et al (2014) <sup>61</sup>	•		•						
Kleinpeter et al (1995)62.			•						
Lee et al (2009)63.		•	•					•	
Martin-Laez et al (2012) <sup>64</sup>			•						
Mayer et al (1993)65.			•			•		•	•
Ohya et al (2016)66.			•						
Pan et al (2014)67.			•						
Righesso et al (2007) <sup>68</sup>	•		•						
Ruetten et al (2008)69.	•	•	•						
Ruetten et al (2009)70.	•	•	•					•	
Sasaoka et al (2006)71.			•						
Schizas et al (2005)72.			•						
Teli et al (2010)73.	•		•						
Ruetten et al (2007)74.			•						
Ruetten et al (2009)75.		•							
Ruetten et al (2008)76.		•						•	
Wang et al (2011)77.		•							
Lee et al (2006)78.	•							•	
Liu et al (2014) <sup>79.</sup>	•								

#### Table 6. Summary of Systematic Reviews of Percutaneous Endoscopic Discectomy Versus Other Discectomy Procedures

Study	Dates	Trials	Participants	N (Range)	Design	Duration
Bai et al (2021) <sup>11.</sup>	To February 2018	14	Patients with lumbar disc herniation who underwent PELD or other surgical procedures	2528 (74-902)	4 RCTs; 10 cohort studies	Not reported
Gadjradj et al (2021) <sup><u>12</u>.</sup>	To April 2020	14	Patients with lumbar disc herniation who underwent PTED or open microdiscectomy	1465 (30-462)	9 RCTs; 5 prospective nonrandomized comparative studies	Follow-up: 3 to 12 months
Xu et al (2020) <u>10.</u>	Search dates not stated; included trials from 2012 to 2018	9	Patients with single-level lumbar disc herniation who underwent PELD or MED for treatment	984 (51-216)	1 Prospective RCTs; 8 Retrospective nonrandomized comparative studies	Follow-up: 1 to > 6 years
Zhou et al (2020) <sup>9</sup> .	To October 2018	12	Patients with lumbar disc herniation who underwent PELD or MED for treatment	2400 (40-915)	4 RCTs; 8 Retrospective nonrandomized comparative studies	Follow-up: 3 to 46 months
Yu et al (2019) <sup>8.</sup>	To August 31, 2018	8	Patients with lumbar disc herniation who underwent PTED or MED procedures and were followed for at least 6 months	805 (51-216)	1 Prospective RCTs; 7 Observational studies	Follow-up: 6 months to 5 years
Shietal (2019) <sup>Z.</sup>	To July 2018	18	Patients with single-level lumbar disc herniation with sciatica who underwent PELD or MED for treatment	2161 (51-273)	8 Prospective studies; 10 Retrospective studies	Follow-up: 3 months to >6 years
Phan et al (2017) <sup>6</sup> .	To February 2016	23	Patients who underwent either an endoscopic or open approach for disc herniation; the endoscopic approach consisted of patients who underwent either FED or MED while the open approach included those who underwent open discectomy or micro-discectomy	28,487 (20-26,612)	10 Prospective RCTs:4 Prospective observational studies;9 Retrospective observational studies	Follow-up: 3 to 104 months
Li et al (2016) <sup>5</sup> .	To January 31, 2015	6	Patients with disc herniation who underwent traditional discectomy surgery or full endoscopic procedures	730 (54-200)	4 RCTs; 2 non-RCTs	Follow-up: 20 to 34 months
Cong et al (2016) <sup><u>4</u>.</sup>	To August 2014	9	Patients who underwent spinal endoscopic or open discectomy for symptomatic lumbar disc herniation	1092 (40-212)	9 RCTs	Follow-up: at least 1 year

FED: full-endoscopic technique discectomy; MED: microendoscopic discectomy; PELD: percutaneous endoscopic lumbar discectomy; PTED: percutaneous transforaminal endoscopic discectomy; RCT: randomized controlled trial

Table 7. Results of Systematic Reviews of Trials of Percutaneous Endoscopic Discectomy Versus Other Discectomy Procedures

Study	Longth of atou	Log poin VAS	Lower back pain VAS	ODI	Querall complication rat-	Beeneration	Desurrence er residue
Study	Length of stay	Leg pain VAS	Lower back pain VAS	001	Overan complication rate	Reoperation	Recurrence of residue
Bai et al (2021)	10	10	10	10	ND		10
Iotal (N)	NK	NR	NK	NK	NK		NK
Pooled effect (95% CI); p value	MD -2.59 (-3.87 to -1.31); <.001	MD 0.00 (-0.10 to 0.10);.991	MD -0.17 (-0.55 to 0.21);.384	MD -0.29 (-1.00 to 0.43);.434	relative risk 0.86 (0.63 to 1.18);.361		relative risk 1.65 (1.08 to 2.52);.021
l <sup>2</sup> (p)	72.1%;.001	0.0%;.996	88.3%; <.001	0.0%;.996	51.5%;.024		26.1%;.220
Gadjradj et al (2021) <sup>12.</sup>							
Total (N)		621 and 152		621 and 152			
Pooled effect (95% CI)		3 to 6 month MD 0.05 (-0.10 to 0.21) 12 month MD 0.11 (-0.30 to 0.53)		3 to 6 month MD -0.09 (-0.24 to 0.07) 12 month MD -0.11 (-0.45 to 0.24)			
l <sup>2</sup> (p)		30%;.23		9%;.83			
Xu et al (2020) <u>10.</u>							
Total (N)	NR	NR	NR	NR	NR	NR	NR
Pooled effect (95% CI); p value	OR -1.041 (-1.493 to -0.583);.000	6 months to 2 years OR -0.138 (-0.384 to 0.108); 270 2 years OR 0.020 (-0.193 to 0.233); 855	6 months to 2 years -0.456 (-0.947 to 0.034);.068 2 years OR -0.856 (-1.488 to -0.224);.008	6 months to 2 years -0.077 (-0.370 to 0.215);.604 2 years OR -0.425 (-0.724 to -0.127);.005	OR 0.972 (0.635 to 1.488);.896	OR 1.136 (0.415 to 3.108);.805	OR 1.306 (0.664 to 2.566); 439
l <sup>2</sup> (p)		53.8%;.090; 6 months to 2 years 4.4%;.351; 2 years	88%;.000; 6 months to 2 years 86.7%;.001; 2 years	75.3%;.000; 6 months to 2 years 52.7%;.121; 2 years			
Zhou et al (2020) <sup>9</sup> .							
Total (N)						787	972
Pooled effect (95% CI); p value						OR 1.77 (1.18 to 2.64);.006	OR 1.60 (1.01 to 2.53);.05
l <sup>2</sup> (p)						0%;.97	0%;.94
Yu et al (2019) <sup>8</sup> .							
Total (N)	707	NR	NR	NR	659		443
Pooled effect (95% CI); p value	MD -1.92 (-2.90 to -0.94); <.001	1 year postop or last follow- up: MD -0.07 (-0.22 to 0.08);.38	1 year postop or last follow-up: MD -0.41 (-0.76 to -0.06);.02	1 year postop or last follow-up: MD -0.27 (-1.71 to 1.16);.71	MD 1.01 (0.60 to 1.69);.98		MD 1.31 (0.54 to 3.17);.54
l <sup>2</sup> (p)	88%				0%		0%
l <sup>2</sup> (p)	96%; <.00001	88%; <.00001	95%; <.00001	55%;.01	0%;.90	0%;.79	0%;.86
Phan et al (2017) <sup>6</sup>							
Total (N)	685	390		303	27,699	995	1081
Pooled effect (95% CI); p value	MD -4.79 (-6.52 to -3.07); <.00001	MD -0.04 (-0.37 to 0.30);.84		MD -1.88 (-4.06 to 0.29);.09	OR 0.77 (0.45 to 1.31);.33	OR 1.46 (0.33 to 6.43);.61	OR 1.12 (0.60 to 2.09);.73
l <sup>2</sup> (p)	99%; <.00001	70%;.003		67%;.03	60%;.004	66%;.004	0%;.97
Li et al (2016) <sup>5.</sup>							
Total (N)	320 (cervical); 410 (lumbar)	410	410	354	730	674	
Pooled effect (95% CI); p value	Cervical: WMD -9.33 (-20.11 to 1.44);.09 Lumbar: WMD -12.16 (-17.24 to -7.09); <.001	At 2 years: -0.58 (-1.46 to 0.29);.19	At 2 years: -1.98 (-6.36 to 2.40);.38	At 2 years: 1.60 (-5.17 to 8.38);.64	RR 0.35 (0.19 to 0.63); <.001	RR 1.02 (0.59 to 1.75);.94	
l <sup>2</sup> (p)	Cervical: 97% Lumbar: 97%	44%;.15	93%; <.001	21%;.28	0%	0%	
Cong et al (2016) <sup>4.</sup>							
Total (N)	NR				NR	NR	NR
Pooled effect (95% CI); p value	WMD -144.45 (-239.54 to -49.37);.003				OR 0.73 (0.34 to 1.57);.41	OR 0.98 (0.60 to 1.61);.93	OR 1.62 (0.84 to 3.12);.15
l <sup>2</sup> (p)	99%				75%	0%	0%
		l			1	1	1

CI: confidence interval; MD: mean difference; NR: not reported; ODI: Oswestry Disability Index; OR: odds ratio; RR: risk ratio; VAS: visual analogue scale; WMD: weighted mean difference

Results from the systematic reviews were fairly consistent with a significantly reduced length of hospitalization observed with endoscopic discectomy and sometimes significant improvements in VAS or ODI, but only at specific time points. Overall, no consistently significant improvement in VAS, ODI, total complication rate, reoperation, or recurrence was observed with endoscopic discectomy versus other interventions. Authors of the systematic reviews noted multiple

limitations including the innate flaws of included studies (i.e., observational designs, a limited number of studies meeting criteria for inclusion, small sample sizes, lack of allocation concealment and blinding), different methodologies contributing to heterogeneity in analyses, loss of usable and sufficient data resulting in difficulty performing accurate analysis of outcomes, and that a majority of the more recently completed studies were completed in China, which may affect the generalizability of the results to other populations.

#### **Randomized Controlled Trials**

A total of 67 trials comparing percutaneous endoscopic discectomy to other discectomy procedures are included in this policy. Sixty-six of these trials were included in at least 1 systematic review (Table 5). More recent RCTs not included in any of the systematic reviews were identified.<sup>80</sup> Results of these trials are similar to those seen in the more comprehensive systematic reviews - percutaneous endoscopic discectomy was associated with a significant reduction in length of stay with no consistent improvement in patient-reported outcome measures such as VAS and ODI. Two of the 3 RCTs evaluated treatment-related morbidities, and reported a reduced incidence of intraoperative and postoperative complications and repeat surgeries with percutaneous endoscopic discectomy. Key characteristics, results, and limitations of these RCTs are summarized in the following tables.

Study	Countries	Sites	Dates	Participants	Interventions
Gadjradj et al 2022 <sup>80,</sup>	Netherlands	4	February 2016 to April 2019	Patients with sciatica caused by lumbar disc herniation	PTED vs microendoscopic discectomy
Ran et al 2021 <sup>81,</sup>	China	1	August 2016 to February 2020	Patients with highly migrated lumbar disc herniation	PELD with computerized tomography navigation vs open discectomy
Wang et al 2019 <sup>82,</sup>	China	1	July 2015 to July 2016	Patients with single-segment lumbar disc herniation with imaging results consistent with symptoms	PTED vs microendoscopic discectomy

#### Table 8. Characteristics of RCTs of Percutaneous Endoscopic Discectomy

PELD: percutaneous endoscopic lumbar discectomy; PTED: percutaneous transforaminal endoscopic discectomy; RCT: randomized controlled trials.

#### Table 9. Results of RCTs of Percutaneous Endoscopic Discectomy

Study	Length of stay (days)	Leg pain VAS	Lower back pain VAS	ODI	SF-36 PCS	Complication rates	Repeat surgery within 1 year
Gadjradj et al 2022 <sup>80,</sup>							
N	420	413	413	413	413	420	420
Pooled effect at 12 months (95% CI)	Median (IQR) PTED: 0 (0 to 0) Microendoscop ic discectomy: 1 (1 to 1)	MD 7.1 (2.8 to 11.3)	MD 6 (2 to 10)	MD 5.3 (3.0 to 7.7)	MD - 2.8 (- 4.1 to - 1.6)	PTED vs microendoscop ic discectomy: Dural tears (n=0 vs 8) Nerve root injury (n=0 vs 1) Wound infection (n=3 vs 0) Cerebrospinal fluid leakage (n=1 vs 0)	PTED vs microendoscop ic discectomy: n=9 (5%) vs 14 (6%)
p-value							
Ran et al 2021 <sup>81,</sup>							
Ν		66				66	
PELD with computerized tomography navigation at 12 months		0.58 ± 0.90				Infection, n=0 Recurrence, n=1	
Open discectomy at 12 months		0.75 ± 0.84				Infection, n=1 Recurrence, n=0	
p-value		.58				>.99	
Ν	90	90	90	90			
PTED	Postoperative: 3.01 ± 0.52	Preoperativ e mean score vs. 6 months after surgery: 7.21 vs. 1.05	Preoperativ e mean score vs 6 months after surgery: 6.40 vs. 1.36	Preope rative mean score vs 6 months after surgery : 58.21% vs. 17.05%			
Microendoscop ic discectomy	Postoperative: 6.68 ± 0.30	Preoperativ e mean score vs. 6 months	Preoperativ e mean score vs 6 months	Preope rative mean score			

		after surgery: 7.09 vs. 0.98	after surgery: 6.34 vs. 1.65	vs 6 months after surgery : 57.17% vs. 16.98%		
p-value	.001	.097	.523	.864		
				2.6		

IQR: interquartile range; MD: mean difference; ODI: Oswestry Disability Index; PELD: percutaneous endoscopic lumbar discectomy; PTED: percutaneous transforaminal endoscopic discectomy; RCT: randomized controlled trials; SF-36 PCS: Short-Form-36 Physical Component Score; VAS: visual analogue scale.

#### Table 10. Study Relevance Limitations of the RCTs of Percutaneous Endoscopic Discectomy

Study	Population <sup>a</sup>	Intervention <sup>b</sup>	Comparator <sup>c</sup>	Outcomes <sup>d</sup>	Duration of Follow-up <sup>e</sup>
Gadjradj et al 2022 <sup>80,</sup>	4. Limited to participants from 3 sites in the Netherlands				
Ran et al 2021 <sup>81,</sup>	4. Limited to participants from single site in China	4.PELD was used with computerized tomography navigation		1. Morbidity- related outcomes such as complications were limited	
Wang et al 2019 <sup>82,</sup>	4. Study population similar to other trials with regard to age, sex; however, included patients from a single Chinese hospital			1. Morbidity- related outcomes such as complication and reoperation rates were not reported	1,2. Outcomes reported only for 6 months of follow-up

PELD: percutaneous endoscopic lumbar discectomy; RCT: randomized controlled trials.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment.

<sup>a</sup> Population key: 1. Intended use population unclear; 2. Study population is unclear; 3. Study population not representative of intended use; 4, Enrolled populations do not reflect relevant diversity; 5. Other.

<sup>b</sup>Intervention key: 1. Not clearly defined; 2. Version used unclear; 3. Delivery not similar intensity as comparator; 4.Not the intervention of interest.

<sup>c</sup>Comparator key: 1. Not clearly defined; 2. Not standard or optimal; 3. Delivery not similar intensity as intervention; 4. Not delivered effectively. <sup>d</sup>Outcomes key: 1. Key health outcomes not addressed; 2. Physiologic measures, not validated surrogates; 3. No CONSORT reporting of harms; 4. Not establish and validated measurements; 5. Clinical significant difference not prespecified; 6. Clinical significant difference not supported.

<sup>e</sup>Follow-Up key: 1. Not sufficient duration for benefit; 2. Not sufficient duration for harms.

#### Table 11. Study Design and Conduct Limitations of the RCTs of Percutaneous Endoscopic Discectomy

Study	Allocation <sup>a</sup>	Blinding <sup>b</sup>	Selective Reporting <sup>c</sup>	Data Completeness <sup>d</sup>	Power <sup>e</sup>	Statistical <sup>f</sup>
Gadjradj et al 2022 <sup>80,</sup>	4. A proportion of patients with a strong preference for PTED who were randomised to open microdiscectomy dropped out of the study after randomization	1,2. Blinding did not occur				
Ran et al 2021 <sup>81,</sup>	3.Allocation concealment unclear	1,2. Blinding did not appear to occur			1. Power calculations not reported	
Wang et al 2019 <sup>82,</sup>	3.Allocation concealment unclear	1,2. Blinding did not appear to occur			1. Power calculations not reported	

PTED: percutaneous transforaminal endoscopic discectomy; RCT: randomized controlled trials.

The study limitations stated in this table are those notable in the current review; this is not a comprehensive gaps assessment. <sup>a</sup>Allocation key: 1. Participants not randomly allocated; 2. Allocation not concealed; 3. Allocation concealment unclear; 4. Inadequate control for selection bias.

<sup>b</sup>Binding key: 1. Not blinded to treatment assignment; 2. Not blinded outcome assessment; 3. Outcome assessed by treating physician.
<sup>c</sup>Selective Reporting key: 1. Not registered; 2. Evidence of selective reporting; 3. Evidence of selective publication.
<sup>d</sup>Data Completeness key: 1. High loss to follow-up or missing data; 2. Inadequate handling of missing data; 3. High number of crossovers; 4. Inadequate handling of crossovers; 5. Inappropriate exclusions; 6. Not intent to treat analysis (per protocol for noninferiority trials).
<sup>e</sup>Power key: 1. Power calculations not reported; 2. Power not calculated for primary outcome; 3. Power not based on clinically important difference.

<sup>1</sup>Statistical key: 1. Analysis is not appropriate for outcome type: (a) continuous; (b) binary; (c) time to event; 2. Analysis is not appropriate for multiple observations per patient; 3. Confidence intervals and/or p values not reported; 4. Comparative treatment effects not calculated.

## **Observational Studies**

Yu et al (2021) published the results of a retrospective multicenter study that followed patients for 2 years after receipt of transforaminal percutaneous endoscopic discectomy (n=632) and microendoscopic discectomy (n=421) for lumbar disc herniation.<sup>8</sup> Mean blood loss (p<.001) and mean duration of hospital stay (p=.018) were significantly less with transforaminal percutaneous endoscopic lumbar discectomy compared to microendoscopic discectomy. Rates of complications, recurrence, and revisions were similar in both groups. Visual analogue pain scores did not differ between groups after the first postoperative day. At 1 month postoperatively there was a significant difference in ODI scores between groups (p=.016) in favor of transforaminal percutaneous endoscopic discectomy, but there was no difference at other time points.

Song et al (2021) published a retrospective single-center study that compared percutaneous endoscopic lumbar discectomy (n=306) and microendoscopic discectomy (n=116) in patients undergoing same day ambulatory surgery for lumbar disc herniation.<sup>37</sup> Mean blood loss and mean duration of hospital stay were significantly less with percutaneous endoscopic lumbar discectomy (both p<.001 compared to microendoscopic discectomy). After 3 years of follow-up, visual analogue pain scores for the back were also significantly lower in the percutaneous

endoscopic lumbar discectomy group compared to the microendoscopic discectomy group (p=.001) but there was no difference between groups in pain scores for the legs (p=.224). Overall recurrence rates (p=.201) and ODI scores (p=.220) were also similar between groups.

A number of observational studies have also assessed the learning curve<sup>83,84,85</sup> and the need for longer follow-up for endoscopic discectomy.<sup>86,87</sup> The largest and longest follow-up to date has been reported by Choi et al (2015), who examined 10,228 patients at their institution who had had percutaneous endoscopic lumbar discectomy over a 12-year period.88 They found that 4.3% of cases required reoperation in the first 6 weeks due to incomplete removal of herniated discs (2.8%), recurrence (0.8%), persistent pain (0.4%), and approach-related pain (0.2%).

## Section Summary: Percutaneous Endoscopic Discectomy

The evidence for percutaneous endoscopic discectomy in individuals who have herniated intervertebral disc(s) includes a number of RCTs and systematic reviews. Many of the more recent RCTs are conducted at institutions within China. There are few reports from the United States. Overall, results from RCTs and systematic reviews reveal a significantly reduced length of hospitalization with endoscopic discectomy and occasionally significant improvements in VAS or ODI, but only at specific time points. No consistently significant improvement in VAS, ODI, total complication rate, reoperation, or recurrence was observed with percutaneous endoscopic discectomy versus other interventions.

## SUMMARY OF EVIDENCE

For individuals who have herniated intervertebral disc(s) who receive automated percutaneous discectomy, the evidence includes randomized controlled trials (RCTs) and systematic reviews of observational studies. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment related morbidity. The published evidence is insufficient to evaluate the impact of automated percutaneous discectomy on net health outcomes. Evidence from small RCTs does not support the use of this procedure. Well-designed and executed RCTs are needed to determine the benefits and risks of this procedure. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have herniated intervertebral disc(s) who receive percutaneous endoscopic discectomy, the evidence includes a number of RCTs and systematic reviews of RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment related morbidity. Many of the RCTs were conducted at a single center in Europe. Some trials have reported outcomes at least as good as traditional approaches with an open incision, while an RCT from a different center in Europe reported a trend toward increased complications and reherniations using an endoscopic approach. There are few reports from the United States. Reporting from a number of moderately large ongoing RCTs is anticipated in the next 2 to 3 years. The evidence is insufficient to determine the effects of the technology on health outcomes.

## **Ongoing and Unpublished Clinical Trials**

Some currently unpublished trials that might influence this review are listed in Table 12.

## Table 12. Summary of Key Trials

Ongoing			
NCT01997086	Percutaneous Transforaminal Endoscopic Discectomy (PTED) vs. Microendoscopic Discectomy (MED) for the treatment of Lumbar Disc Herniation: A Prospective Randomized Controlled Study		Aug 2023
Unpublished			
NCT02742311	EuroPainClinics® Study V Prospective Observational Study (EPCSV)	500	Dec 2021

NCT: national clinical trial

<sup>a</sup> Denotes industry-sponsored or cosponsored trial

#### SUPPLEMENTAL INFORMATION

# Clinical Input Obtained Through Physician Specialty Societies and Academic Medical Centers

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

#### 2018 Input

In response to requests, clinical input on automated percutaneous discectomy and percutaneous endoscopic discectomy for herniated intervertebral disc(s) was received from 3 respondents, including 2 specialty society-level responses and including physicians with academic medical center affiliation, while this policy was under review in 2018.

#### 2013 Input

In response to requests, input was received from 4 physician specialty societies and 3 academic medical centers while this policy was under review in 2013. Overall, input agreed that percutaneous and endoscopic discectomy are investigational. Most reviewers considered discectomy with tubular retractors to be a variant of open discectomy, with the only difference being the type of retraction used.

#### PRACTICE GUIDELINES AND POSITION STATEMENTS

#### National Institute for Health and Clinical Excellence

The National Institute for Health and Clinical Excellence (NICE) published guidance in 2005 on automated percutaneous mechanical lumber discectomy, indicating that there is limited evidence of efficacy based on uncontrolled case series of heterogeneous groups of patients, and evidence from small randomized controlled trials shows conflicting results.<sup>89</sup> The guidance states that in view of uncertainty about the efficacy of the procedure, it should not be done without special arrangements for consent and for audit or research. The guidance was considered for update in 2009, but failed review criteria; the 2005 guidance is therefore considered current.

A NICE guideline on percutaneous transforaminal endoscopic lumbar discectomy for sciatica went in development in March 2016.<sup>90</sup> The guidance stated that current evidence is adequate to support the use of percutaneous transforaminal endoscopic lumbar discectomy for sciatica.

Choice of operative procedure (open discectomy, microdiscectomy, or percutaneous endoscopic approaches) may be influenced by symptoms, location and size of prolapsed disc.

A NICE guidance on percutaneous interlaminar endoscopic lumbar discectomy for sciatica was also published in 2016.<sup>91</sup> The guidance stated that current evidence is adequate to support the use of percutaneous interlaminar endoscopic lumbar discectomy for sciatica. Choice of operative procedure (open discectomy, microdiscectomy, or percutaneous endoscopic approaches) may be influenced by symptoms, and location and size of prolapsed disc.

#### **American Society of Interventional Pain Physicians**

The 2013 guideline update from the American Society of Interventional Pain Physicians states that the evidence for percutaneous disc decompression with Dekompressor is limited.<sup>3</sup> There were no recommended indications for DeKompressor.

#### North American Spine Society

In 2014, the North American Spine Society published clinical guidelines on the diagnosis and treatment of lumbar disc herniation.<sup>92</sup> Table 13 summarizes recommendations specific to percutaneous endoscopic discectomy and automated percutaneous discectomy.

#### Table 13. NASS Recommendations for Lumbar Disc Herniation With Radiculopathy

Recommendations	Grade or LOE <sup>a</sup>
Endoscopic percutaneous discectomy is suggested for carefully selected patients to reduce early postoperative disability and reduce opioid use compared with open discectomy	В
There is insufficient evidence to make a recommendation for or against the use of automated percutaneous discectomy compared with open discectomy	I
Endoscopic percutaneous discectomy may be considered for treatment	С
Automated percutaneous discectomy may be considered for treatment	С
Patients undergoing percutaneous endoscopic discectomy experience better outcomes if <40 years and symptom duration <3 months	II

LOE: level of evidence; NASS: North American Spine Society

<sup>a</sup> Grade B: fair evidence (level II or III studies with consistent findings; grade C: poor quality evidence (Level IV or V studies). Level of evidence II: lesser quality randomized controlled trial (e.g., <80% follow-up, no blinding, or improper randomization), prospective comparative study, systematic review of level II studies or level I studies with inconsistent results; level of evidence III: case control, retrospective, systematic review of level III studies; level of evidence IV: case series; level of evidence V: expert opinion.

#### **American Pain Society**

The 2009 clinical practice guidelines from the American Pain Society found insufficient evidence to evaluate alternative surgical methods to standard open discectomy and microdiscectomy, including laser or endoscopic-assisted techniques, various percutaneous techniques, coblation nucleoplasty, or the Disc Dekompressor.<sup>93</sup>

# **Government Regulations**

#### National/ Local:

There are no national or local coverage determinations on this topic. 62380 is not on the Medicare Fee schedule. 62287 can be used for TIPS procedures and is not covered for that indication; however 62287 is payable for procedures that are not considered TIPS procedures.

National Coverage Determination, Publication 100-3, Manual Section Number 150.11. Percutaneous thermal intradiscal procedures (TIPs) involve the insertion a catheter(s)/probe(s) into the spinal disc under fluoroscopic guidance in order to produce, or apply, heat and/or disruption within the disc to relieve low back pain.

Although not meant to be a complete list, TIPs are commonly identified as

- Intradiscal electrothermal therapy (IDET);
- Intradiscal thermal annuloplasty (IDTA);
- Percutaneous intradiscal radiofrequency thermocoagulation (PIRFT);
- Radiofrequency annuloplasty (RA);
- Intradiscal biacuplasty (IDB);
- Percutaneous (or plasma) disc decompression (PDD) or ablation; or
- Targeted disc decompression (TDD).

CR 6291 announces that CMS has concluded that the evidence does not demonstrate that TIPs improve health outcomes; and has therefore determined that **TIPs are not reasonable and necessary** for the treatment of low back pain.

(The above Medicare information is current as of the review date for this policy. However, the coverage issues and policies maintained by the Centers for Medicare & Medicare Services [CMS, formerly HCFA] are updated and/or revised periodically. Therefore, the most current CMS information may not be contained in this document. For the most current information, the reader should contact an official Medicare source.

# **Related Policies**

• N/A

## References

- Lewis RA, Williams NH, Sutton AJ, et al. Comparative clinical effectiveness of management strategies for sciatica: systematic review and network meta-analyses. Spine J. Jun 01 2015; 15(6): 1461-77. PMID 24412033
- 2. Haines SJ, Jordan N, Boen JR, et al. Discectomy strategies for lumbar disc herniation: results of the LAPDOG trial. J Clin Neurosci. Jul 2002; 9(4): 411-7. PMID 12217670
- 3. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II: guidance and recommendations. Pain Physician. Apr 2013; 16(2 Suppl): S49-283. PMID 23615883
- Cong L, Zhu Y, Tu G. A meta-analysis of endoscopic discectomy versus open discectomy for symptomatic lumbar disk herniation. Eur Spine J. Jan 2016; 25(1): 134-143. PMID 25632840
- Li XC, Zhong CF, Deng GB, et al. Full-Endoscopic Procedures Versus Traditional Discectomy Surgery for Discectomy: A Systematic Review and Meta-analysis of Current Global Clinical Trials. Pain Physician. Mar 2016; 19(3): 103-18. PMID 27008284

- Phan K, Xu J, Schultz K, et al. Full-endoscopic versus micro-endoscopic and open discectomy: A systematic review and meta-analysis of outcomes and complications. Clin Neurol Neurosurg. Mar 2017; 154: 1-12. PMID 28086154
- 7. Shi R, Wang F, Hong X, et al. Comparison of percutaneous endoscopic lumbar discectomy versus microendoscopic discectomy for the treatment of lumbar disc herniation: a metaanalysis. Int Orthop. Apr 2019; 43(4): 923-937. PMID 30547214
- Yu P, Qiang H, Zhou J, et al. Percutaneous Transforaminal Endoscopic Discectomy versus Micro-Endoscopic Discectomy for Lumbar Disc Herniation. Med Sci Monit. Mar 30 2019; 25: 2320-2328. PMID 30927349
- Zhao XM, Yuan QL, Liu L, et al. Is It Possible to Replace Microendoscopic Discectomy with Percutaneous Transforaminal Discectomy for Treatment of Lumbar Disc Herniation? A Meta-Analysis Based on Recurrence and Revision Rate. J Korean Neurosurg Soc. Jul 2020; 63(4): 477-486. PMID 32380585
- 10. Xu J, Li Y, Wang B, et al. Minimum 2-Year Efficacy of Percutaneous Endoscopic Lumbar Discectomy versus Microendoscopic Discectomy: A Meta-Analysis. World Neurosurg. Jun 2020; 138: 19-26. PMID 32109644
- 11. Bai X, Lian Y, Wang J, et al. Percutaneous endoscopic lumbar discectomy compared with other surgeries for lumbar disc herniation: A meta-analysis. Medicine (Baltimore). Mar 05 2021; 100(9): e24747. PMID 33655938
- 12. Gadjradj PS, Harhangi BS, Amelink J, et al. Percutaneous Transforaminal Endoscopic Discectomy Versus Open Microdiscectomy for Lumbar Disc Herniation: A Systematic Review and Meta-analysis. Spine (Phila Pa 1976). Apr 15 2021; 46(8): 538-549. PMID 33290374
- 13. Akcakaya MO, Yorukoglu AG, Aydoseli A, et al. Serum creatine phosphokinase levels as an indicator of muscle injury following lumbar disc surgery: Comparison of fully endoscopic discectomy and microdiscectomy. Clin Neurol Neurosurg. Jun 2016; 145: 74-8. PMID 27101087
- 14. Choi KC, Shim HK, Hwang JS, et al. Comparison of Surgical Invasiveness Between Microdiscectomy and 3 Different Endoscopic Discectomy Techniques for Lumbar Disc Herniation. World Neurosurg. Aug 2018; 116: e750-e758. PMID 29787880
- 15. Dai HJ, Zhang X, Wang LT, et al. The effect of percutaneous transforaminal endoscopic discectomy (PTED) on serum inflammatory factors and pain in patients with lumbar disc herniation after surgery. Int J Clin Exp Med 2020;13:597603.
- 16. Krappel FA, Schmitz R, Bauer E, et al. Open or endoscopic nucleotomy?. Orthopadische Praxis 2001;37:1649.
- 17. Tacconi L, Giordan E. Endoscopic transforaminal discectomy vs. far lateral discectomy for extraforaminal disc protrusions: our experience. NeuroQuantology 2019;17:1822.
- Tacconi L, Signorelli F, Giordan E. Is Full Endoscopic Lumbar Discectomy Less Invasive Than Conventional Surgery? A Randomized MRI Study. World Neurosurg. Jun 2020; 138: e867-e875. PMID 32251813
- Tao XZ, Jing L, Li JH. Therapeutic effect of transforaminal endoscopic spine system in the treatment of prolapse of lumbar intervertebral disc. Eur Rev Med Pharmacol Sci. Jul 2018; 22(1 Suppl): 103-110. PMID 30004561
- 20. Wang H, Song Y, Cai L. Effect of percutaneous transforaminal lumbar spine endoscopic discectomy on lumbar disc herniation and its influence on indexes of oxidative stress. Biomed Res 2017;28:.
- 21. Xu G, Zhang C, Zhu K, et al. Endoscopic removal of nucleus pulposus of intervertebral disc on lumbar intervertebral disc protrusion and the influence on inflammatory factors and immune function. Exp Ther Med. Jan 2020; 19(1): 301-307. PMID 31853303

- 22. Ahn SS, Kim SH, Kim DW, et al. Comparison of Outcomes of Percutaneous Endoscopic Lumbar Discectomy and Open Lumbar Microdiscectomy for Young Adults: A Retrospective Matched Cohort Study. World Neurosurg. Feb 2016; 86: 250-8. PMID 26409086
- 23. Chang F, Zhang T, Gao G, et al. Therapeutic effect of percutaneous endoscopic lumbar discectomy on lumbar disc herniation and its effect on oxidative stress in patients with lumbar disc herniation. Exp Ther Med. Jan 2018; 15(1): 295-299. PMID 29250152
- 24. Liu C, Zhou Y. Percutaneous Endoscopic Lumbar Diskectomy and Minimally Invasive Transforaminal Lumbar Interbody Fusion for Recurrent Lumbar Disk Herniation. World Neurosurg. Feb 2017; 98: 14-20. PMID 27773858
- 25. Pan Z, Ha Y, Yi S, et al. Efficacy of Transforaminal Endoscopic Spine System (TESSYS) Technique in Treating Lumbar Disc Herniation. Med Sci Monit. Feb 18 2016; 22: 530-9. PMID 26887645
- 26. Yao Y, Zhang H, Wu J, et al. Comparison of Three Minimally Invasive Spine Surgery Methods for Revision Surgery for Recurrent Herniation After Percutaneous Endoscopic Lumbar Discectomy. World Neurosurg. Apr 2017; 100: 641-647.e1. PMID 28153616
- 27. Yao Y, Zhang H, Wu J, et al. Minimally Invasive Transforaminal Lumbar Interbody Fusion Versus Percutaneous Endoscopic Lumbar Discectomy: Revision Surgery for Recurrent Herniation After Microendoscopic Discectomy. World Neurosurg. Mar 2017; 99: 89-95. PMID 27919762
- 28. Gibson JNA, Subramanian AS, Scott CEH. A randomised controlled trial of transforaminal endoscopic discectomy vs microdiscectomy. Eur Spine J. Mar 2017; 26(3): 847-856. PMID 27885470
- 29. Hsu HT, Chang SJ, Yang SS, et al. Learning curve of full-endoscopic lumbar discectomy. Eur Spine J. Apr 2013; 22(4): 727-33. PMID 23076645
- 30. Kim MJ, Lee SH, Jung ES, et al. Targeted percutaneous transforaminal endoscopic diskectomy in 295 patients: comparison with results of microscopic diskectomy. Surg Neurol. Dec 2007; 68(6): 623-631. PMID 18053857
- 31. Qu JX, Li QZ, Chen M : Comparative study of PTED and MED for monosegmentnlumbar disc herniation. Chin J Bone Joint Inj 32 : 70-71,2017
- 32. Wang H, Cheng J, Xiao H, et al. Adolescent lumbar disc herniation: experience from a large minimally invasive treatment centre for lumbar degenerative disease in Chongqing, China. Clin Neurol Neurosurg. Aug 2013; 115(8): 1415-9. PMID 23419406
- 33. Zhao W, Li CQ, Zhou Y, Wang J, Zheng WJ : Surgical treatment of thenlumbar disc herniated discs using transforaminal endoscopic surgery system. Orthop J China 20 : 1191-1195, 2012
- 34. Yoon SM, Ahn SS, Kim KH, et al. Comparative Study of the Outcomes of Percutaneous Endoscopic Lumbar Discectomy and Microscopic Lumbar Discectomy Using the Tubular Retractor System Based on the VAS, ODI, and SF-36. Korean J Spine. Sep 2012; 9(3): 215-22. PMID 25983818
- 35. Li M, Yang H, Yang Q. Full-Endoscopic Technique Discectomy Versus Microendoscopic Discectomy for the Surgical Treatment of Lumbar Disc Herniation. Pain Physician. Jul-Aug 2015; 18(4): 359-63. PMID 26218939
- 36. Sinkemani A, Hong X, Gao ZX, et al. Outcomes of Microendoscopic Discectomy and Percutaneous Transforaminal Endoscopic Discectomy for the Treatment of Lumbar Disc Herniation: A Comparative Retrospective Study. Asian Spine J. Dec 2015; 9(6): 833-40. PMID 26713113
- 37. Song HP, Sheng HF, Xu WX. A case-control study on the treatment of protrusion of lumbar intervertebral disc through PELD and MED. Exp Ther Med. Oct 2017; 14(4): 3708-3712. PMID 29042967

- 38. Tu Z, Li YW, Wang B, et al. Clinical Outcome of Full-endoscopic Interlaminar Discectomy for Single-level Lumbar Disc Herniation: A Minimum of 5-year Follow-up. Pain Physician. Mar 2017; 20(3): E425-E430. PMID 28339442
- 39. Liu X, Yuan S, Tian Y, et al. Comparison of percutaneous endoscopic transforaminal discectomy, microendoscopic discectomy, and microdiscectomy for symptomatic lumbar disc herniation: minimum 2-year follow-up results. J Neurosurg Spine. Mar 2018; 28(3): 317-325. PMID 29303471
- 40. Li H, Jiang C, Mu X, et al. Comparison of MED and PELD in the Treatment of Adolescent Lumbar Disc Herniation: A 5-Year Retrospective Follow-Up. World Neurosurg. Apr 2018; 112: e255-e260. PMID 29325949
- 41. Abudurexiti T, Qi L, Muheremu A, et al. Micro-endoscopic discectomy versus percutaneous endoscopic surgery for lumbar disk herniation. J Int Med Res. Sep 2018; 46(9): 3910-3917. PMID 29900752
- 42. Chen Z, Zhang L, Dong J, et al. Percutaneous transforaminal endoscopic discectomy compared with microendoscopic discectomy for lumbar disc herniation: 1-year results of an ongoing randomized controlled trial. J Neurosurg Spine. Mar 2018; 28(3): 300-310. PMID 29303469
- 43. Liu T, Zhou Y, Wang J, et al. Clinical efficacy of three different minimally invasive procedures for far lateral lumbar disc herniation. Chin Med J (Engl). Mar 2012; 125(6): 1082-8. PMID 22613535
- 44. Wu XC, Zhou Y, Li CQ. Percutaneous tranforaminal endoscopic discectomy versus microendoscopic discectomy for lumbar disc herniation: a prospective randomized controlled study. J Third Mil Med Univ. 2009;31(9):843-846.
- 45. Yang L, Liao XQ, Zhao XJ, et al. Comparison of surgical outcomes between percutaneous transforaminal endoscopic discectomy and micro-endoscopic discectomy for lumbar disc herniation. China J Endosc. 2015;21(9):962-965
- 46. Duan XF, Jin W, Chen JJ, et al. Contrast observation of comparing microendoscopic discectomy with percutaneous endoscopic lumbar discectomy for the treatment of simple lumbar disc herniation. Chin J Clin. 2016;10(1):144-147
- 47. Zhao XW, Han K, Ji ZW, et al. Comparison of efficacy between microendoscopic discectomy and percutaneous endoscopic lumbar discectomy for treatment of lumbar disc herniation. Prog Mod Biomed. 2016;16(23):4454-4457
- 48. Ding YZ, Hu JN, Zhou Y, et al. Study on the effect contrast between microendoscopic discectomy and percutaneous endoscopic lumbar discectomy using TESSYS technique for the treatment of lumbar disc herniation. J Cervicodynia & Lumbodynia. 2017;38(5):492-493
- 49. Li ZY, Guo PG, Han D, et al. Analysis of curative effects and prognosis in different procedures of discectomy for patients with lumbar disc herniation. J Clin Med Pract. 2017;21(15):149-150,158
- 50. Liu HP, Hao DJ, Wang XD, et al. Comparison of two surgeries in treatment of lumbar disc herniation. Chin J Pain Med. 2017;23(6):438-442
- 51. Luo DK, Zhou NX, Zhao HW, et al. Clinical effectiveness of minimally invasive treatment for lumbar disc herniation. Orthopaedics. 2017;8(6):439-444
- 52. Qu JX, Li QZ, Chem M, et al. Comparison of the efficacies between percutaneous transforaminal endoscopic discectomy and microendoscopic discectomy for the treatment of single-segmental lumbar disc herniation. Chin J Bone Jt Inj. 2017;32(1):70-71
- 53. Chen Q, Qin L, Li MW, et al. Comparison of the therapeutic effect of percutaneous transforaminal endoscopic discectomy and posterior discectomy on senile single segmental lumbar disc herniation. Chin J Front Med Sci. 2018;10(2):60-64

- 54. Wu YM, Bai M, Yin HP, et al. Comparison of the efficacies between two kinds of minimally invasive procedures for the treatment of simple lumbar disc herniation. J Pract Orthop. 2018;24(4):357-360
- 55. Belykh E, Giers MB, Preul MC, et al. Prospective Comparison of Microsurgical, Tubular-Based Endoscopic, and Endoscopically Assisted Diskectomies: Clinical Effectiveness and Complications in Railway Workers. World Neurosurg. Jun 2016; 90: 273-280. PMID 26898494
- 56. Chen HC, Lee CH, Wei L, et al. Comparison of percutaneous endoscopic lumbar discectomy and open lumbar surgery for adjacent segment degeneration and recurrent disc herniation. Neurol Res Int. 2015; 2015: 791943. PMID 25861474
- 57. Choi KC, Kim JS, Park CK. Percutaneous Endoscopic Lumbar Discectomy as an Alternative to Open Lumbar Microdiscectomy for Large Lumbar Disc Herniation. Pain Physician. Feb 2016; 19(2): E291-300. PMID 26815256
- 58. Garg B, Nagraja UB, Jayaswal A. Microendoscopic versus open discectomy for lumbar disc herniation: a prospective randomised study. J Orthop Surg (Hong Kong). Apr 2011; 19(1): 30-4. PMID 21519072
- 59. Hermantin FU, Peters T, Quartararo L, et al. A prospective, randomized study comparing the results of open discectomy with those of video-assisted arthroscopic microdiscectomy. J Bone Joint Surg Am. Jul 1999; 81(7): 958-65. PMID 10428127
- 60. Huang TJ, Hsu RW, Li YY, et al. Less systemic cytokine response in patients following microendoscopic versus open lumbar discectomy. J Orthop Res. Mar 2005; 23(2): 406-11. PMID 15734255
- Hussein M, Abdeldayem A, Mattar MM. Surgical technique and effectiveness of microendoscopic discectomy for large uncontained lumbar disc herniations: a prospective, randomized, controlled study with 8 years of follow-up. Eur Spine J. Sep 2014; 23(9): 1992-9. PMID 24736930
- 62. Kleinpeter G, Markowitsch MM, Bock F. Percutaneous endoscopic lumbar discectomy: minimally invasive, but perhaps only minimally useful?. Surg Neurol. Jun 1995; 43(6): 534-9; discussion 540-1. PMID 7482230
- 63. Lee DY, Shim CS, Ahn Y, et al. Comparison of percutaneous endoscopic lumbar discectomy and open lumbar microdiscectomy for recurrent disc herniation. J Korean Neurosurg Soc. Dec 2009; 46(6): 515-21. PMID 20062565
- 64. Martin-Laez R, Martinez-Agueros JA, Suarez-Fernandez D, et al. Complications of endoscopic microdiscectomy using the EASYGO! system: is there any difference with conventional discectomy during the learning-curve period?. Acta Neurochir (Wien). Jun 2012; 154(6): 1023-32. PMID 22446750
- 65. Mayer HM, Brock M. Percutaneous endoscopic discectomy: surgical technique and preliminary results compared to microsurgical discectomy. J Neurosurg. Feb 1993; 78(2): 216-25. PMID 8267686
- 66. Ohya J, Oshima Y, Chikuda H, et al. Does the microendoscopic technique reduce mortality and major complications in patients undergoing lumbar discectomy? A propensity scorematched analysis using a nationwide administrative database. Neurosurg Focus. Feb 2016; 40(2): E5. PMID 26828886
- 67. Pan L, Zhang P, Yin Q. Comparison of tissue damages caused by endoscopic lumbar discectomy and traditional lumbar discectomy: a randomised controlled trial. Int J Surg. 2014; 12(5): 534-7. PMID 24583364
- 68. Righesso O, Falavigna A, Avanzi O. Comparison of open discectomy with microendoscopic discectomy in lumbar disc herniations: results of a randomized controlled trial. Neurosurgery. Sep 2007; 61(3): 545-9; discussion 549. PMID 17881967

- 69. Ruetten S, Komp M, Merk H, et al. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study. Spine (Phila Pa 1976). Apr 20 2008; 33(9): 931-9. PMID 18427312
- 70. Ruetten S, Komp M, Merk H, et al. Recurrent lumbar disc herniation after conventional discectomy: a prospective, randomized study comparing full-endoscopic interlaminar and transforaminal versus microsurgical revision. J Spinal Disord Tech. Apr 2009; 22(2): 122-9. PMID 19342934
- 71. Sasaoka R, Nakamura H, Konishi S, et al. Objective assessment of reduced invasiveness in MED. Compared with conventional one-level laminotomy. Eur Spine J. May 2006; 15(5): 577-82. PMID 15926058
- 72. Schizas C, Tsiridis E, Saksena J. Microendoscopic discectomy compared with standard microsurgical discectomy for treatment of uncontained or large contained disc herniations. Neurosurgery. Oct 2005; 57(4 Suppl): 357-60; discussion 357-60. PMID 16234685
- 73. Teli M, Lovi A, Brayda-Bruno M, et al. Higher risk of dural tears and recurrent herniation with lumbar micro-endoscopic discectomy. Eur Spine J. Mar 2010; 19(3): 443-50. PMID 20127495
- 74. Ruetten S, Komp M, Merk H, et al. Use of newly developed instruments and endoscopes: full-endoscopic resection of lumbar disc herniations via the interlaminar and lateral transforaminal approach. J Neurosurg Spine. Jun 2007; 6(6): 521-30. PMID 17561740
- 75. Ruetten S, Komp M, Merk H, et al. Full-endoscopic anterior decompression versus conventional anterior decompression and fusion in cervical disc herniations. Int Orthop. Dec 2009; 33(6): 1677-82. PMID 19015851
- 76. Ruetten S, Komp M, Merk H, et al. Full-endoscopic cervical posterior foraminotomy for the operation of lateral disc herniations using 5.9-mm endoscopes: a prospective, randomized, controlled study. Spine (Phila Pa 1976). Apr 20 2008; 33(9): 940-8. PMID 18427313
- 77. Wang B, Lu GH, Li J, et al. [Contrast study of full-endoscopic interlaminar approach for the surgical treatment of lumbar disc herniation]. Zhonghua Wai Ke Za Zhi. Jan 01 2011; 49(1): 74-8. PMID 21418844
- 78. Lee SH, Chung SE, Ahn Y, et al. Comparative radiologic evaluation of percutaneous endoscopic lumbar discectomy and open microdiscectomy: a matched cohort analysis. Mt Sinai J Med. Sep 2006; 73(5): 795-801. PMID 17008941
- 79. Liu JL, Zhen WX, Gao GY, et al. A prospective and controlled study of percutaneous transforaminal endoscopic discectomy versus fenestration discectomy for lumbar disc herniation. Chin J Bone Joint. 2014;3:245-250
- 80. Gadjradj PS, Rubinstein SM, Peul WC, et al. Full endoscopic versus open discectomy for sciatica: randomised controlled non-inferiority trial. BMJ. Feb 21 2022; 376: e065846. PMID 35190388
- 81. Ran B, Wei J, Yang J, et al. Quantitative Evaluation of the Trauma of CT Navigation PELD and OD in the Treatment of HLDH: A Randomized, Controlled Study. Pain Physician. Jul 2021; 24(4): E433-E441. PMID 34213868
- 82. Wang F, Guo D, Sun T, et al. A comparative study on short-term therapeutic effects of percutaneous transforaminal endoscopic discectomy and microendoscopic discectomy on lumbar disc herniation. Pak J Med Sci. Mar-Apr 2019; 35(2): 426-431. PMID 31086527
- 83. Lee DY, Lee SH. Learning curve for percutaneous endoscopic lumbar discectomy. Neurol Med Chir (Tokyo). Sep 2008; 48(9): 383-8; discussion 388-9. PMID 18812679
- 84. Wang B, Lu G, Patel AA, et al. An evaluation of the learning curve for a complex surgical technique: the full endoscopic interlaminar approach for lumbar disc herniations. Spine J. Feb 2011; 11(2): 122-30. PMID 21296295

- 85. Tenenbaum S, Arzi H, Herman A, et al. Percutaneous Posterolateral Transforaminal Endoscopic Discectomy: Clinical Outcome, Complications, and Learning Curve Evaluation. Surg Technol Int. Dec 2011; 21: 278-83. PMID 22505002
- 86. Casal-Moro R, Castro-Menendez M, Hernandez-Blanco M, et al. Long-term outcome after microendoscopic diskectomy for lumbar disk herniation: a prospective clinical study with a 5-year follow-up. Neurosurgery. Jun 2011; 68(6): 1568-75; discussion 1575. PMID 21311384
- Wang M, Zhou Y, Wang J, et al. A 10-year follow-up study on long-term clinical outcomes of lumbar microendoscopic discectomy. J Neurol Surg A Cent Eur Neurosurg. Aug 2012; 73(4): 195-8. PMID 22825836
- 88. Choi KC, Lee JH, Kim JS, et al. Unsuccessful percutaneous endoscopic lumbar discectomy: a single-center experience of 10,228 cases. Neurosurgery. Apr 2015; 76(4): 372-80; discussion 380-1; quiz 381. PMID 25599214
- 89. National Institute for Health and Care Excellence (NICE). Automated percutaneous mechanical lumbar discectomy-guidance [IPG141]. 2005; http://guidance.nice.org.uk/IPG141/Guidance/pdf/English. Accessed July 2022
- 90. National Institute for Health and Care Excellence (NICE). Percutaneous interlaminar endoscopic lumbar discectomy for sciatica [IPG555]. 2016; https://www.nice.org.uk/guidance/ipg555. Accessed July 2022
- 91. National Institute for Health and Care Excellence (NICE). Percutaneous transforaminal endoscopic lumbar discectomy for sciatica [IPG556]. 2016; https://www.nice.org.uk/guidance/ipg556. Accessed July 2022.
- 92. Kreiner DS, Hwang SW, Easa JE, et al. An evidence-based clinical guideline for the diagnosis and treatment of lumbar disc herniation with radiculopathy. Spine J. Jan 2014; 14(1): 180-91. PMID 24239490
- 93. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. Spine (Phila Pa 1976). May 01 2009; 34(10): 1066-77. PMID 19363457
- 95. Blue Cross Blue Shield Association. Automated Percutaneous and Endoscopic Discectomy. Medical Policy Reference Manual. MPRM 7.01.18. Last updated July 2022.
- 96. HAYES Directory Assessment. Automated Percutaneous Lumbar Discectomy. Lansdale, PA: HAYES, Inc. December 4, 2014. Last updated November 2016. Archived January 2019.
- 97 HAYES Health Technology Brief. Arthroscopic Microdiscectomy for Lumbar Disc Herniation. Lansdale, PA: HAYES, Inc. December 28, 2011, updated January 6, 2014. Archived January 2015.
- 98. HAYES Search and Summary. Arthroscopic Microdiscectomy for Lumbar Disc Herniation. Lansdale, PA: HAYES, Inc. published September 8, 2016.
- HAYES Health Technology Brief. METRx Microscope System (Medtronic Sofamor Danek Inc.) for Microdiscectomy for Lumbar Disc Herniation. Lansdale, PA: HAYES, Inc. December 24, 2012, updated December 2014. Archived January 2016; Outdated.
- 100. HAYES Health Technology Brief. Percutaneous endoscopic lumbar discectomy for primary lumbar disc herniation. Lansdale, PA: HAYES, Inc. Published March 02, 2017. Last updated February 2019.
- 101. Hayes Health Technology Brief. Percutaneous endoscopic lumbar discectomy for recurrent lumbar disc herniation. Lansdale, PA: HAYES, Inc. Published March 23, 2017.

The articles reviewed in this research include those obtained in an Internet based literature search for relevant medical references through July 2022, the date the research was completed.

# **Medical Policy History**

Policy Effective Date	Comments
1/1/2022	Medical policy established
11/1/2022	Rationale section updated, added references 80-82. No change in policy status.

Next Review Date: 3<sup>rd</sup> Qtr. 2023