

2016 Procedures Criteria

PATIENT:	Name	DOB	ID#	GROUP#
	Facility		Service Date	
PROVIDER:	Name		Fax#	Phone#
	Signature		Date	NPI/ID#

ICD-9:

ICD-10:

CPT®:

Subset: Vertebroplasty or Kyphoplasty^(1, 2, 3, 4)**Requested Service:** Vertebroplasty**Age:** Age ≥ 18**INSTRUCTIONS:** Choose one of the following options and continue to the appropriate section

10. Malignancy related compression fracture
20. Osteoporotic vertebral compression fracture
30. Vertebral hemangioma

 10. Malignancy related compression fracture

1. Choose all that apply:

- A) Back pain interferes with ADLs
- B) No neurologic deficits⁽⁵⁾
- C) Compression fracture by imaging
- D) Other clinical information (add comment)

- If the number of options selected is 3 and option D not selected, then go to question 2
- No other options lead to the requested service

2. Choose all that apply:^(6, 7)

- A) Malignancy by biopsy
- B) Evidence of malignancy by history
- C) Evidence of malignancy by current radiology report
- D) Other clinical information (add comment)

- If 1 or more options A, B or C selected and option D not selected, then go to question 3
- No other options lead to the requested service

3. Choose all that apply:⁽⁸⁾

InterQual® criteria (IQ) is confidential and proprietary information and is being provided to you solely as it pertains to the information requested. IQ may contain advanced clinical knowledge which we recommend you discuss with your physician upon disclosure to you. Use permitted by and subject to license with McKesson Corporation and/or one of its subsidiaries. IQ reflects clinical interpretations and analyses and cannot alone either (a) resolve medical ambiguities of particular situations; or (b) provide the sole basis for definitive decisions. IQ is intended solely for use as screening guidelines with respect to medical appropriateness of healthcare services. All ultimate care decisions are strictly and solely the obligation and responsibility of your health care provider. InterQual® and CareEnhance® Review Manager © 2016 McKesson Corporation and/or one of its subsidiaries. All Rights Reserved. CPT only © 2015 American Medical Association. All Rights Reserved.

10. Malignancy related compression fracture (*Continued...*)

- A) No septicemia
- B) No active osteomyelitis of the target vertebra
- C) No coagulopathy or coagulopathy treated
- D) No known allergy to bone cement or opacification agent
- E) Other clinical information (add comment)

- If the number of options selected is 4 and option E not selected, then the rule is satisfied; you may stop here (*Outpatient*)
- No other options lead to the requested service

 20. Osteoporotic vertebral compression fracture

1. Choose all that apply:

- A) Osteoporosis by history
- B) Back pain interferes with ADLs
- C) No neurologic deficits⁽⁵⁾
- D) Osteoporotic compression fracture by imaging
- E) Other clinical information (add comment)

- If the number of options selected is 4 and option E not selected, then go to question 2
- No other options lead to the requested service

2. Treatment within last year, Choose all that apply:⁽⁹⁾

- A) NSAIDs or acetaminophen \geq 3 weeks⁽¹⁰⁾
- B) Back brace \geq 4 weeks or back brace contraindicated or not tolerated⁽¹¹⁾
- C) Activity modification \geq 6 weeks⁽¹²⁾
- D) Other clinical information (add comment)

- If the number of options selected is 3 and option D not selected, then go to question 3
- No other options lead to the requested service

3. Continued pain after treatment

- Yes
- No

- If option Yes selected, then go to question 4
- No other options lead to the requested service

4. Choose all that apply:⁽⁸⁾

20. Osteoporotic vertebral compression fracture (*Continued...*)

- A) No septicemia
- B) No active osteomyelitis of the target vertebra
- C) No coagulopathy or coagulopathy treated
- D) No known allergy to bone cement or opacification agent
- E) Other clinical information (add comment)

- If the number of options selected is 4 and option E not selected, then the rule is satisfied; you may stop here (*Outpatient*)
- No other options lead to the requested service

 30. Vertebral hemangioma

1. Choose all that apply:

- A) Back pain interferes with ADLs
- B) No neurologic deficits⁽⁵⁾
- C) Pathologic fracture by imaging⁽¹³⁾
- D) Other clinical information (add comment)

- If the number of options selected is 3 and option D not selected, then go to question 2
- No other options lead to the requested service

2. Treatment within last year, Choose all that apply:

- A) NSAIDs or acetaminophen \geq 3 weeks⁽¹⁰⁾
- B) Back brace \geq 4 weeks OR back brace contraindicated or not tolerated⁽¹¹⁾
- C) Activity modification \geq 6 weeks⁽¹²⁾
- D) Other clinical information (add comment)

- If the number of options selected is 3 and option D not selected, then go to question 3
- No other options lead to the requested service

3. Continued pain after treatment

- Yes
- No

- If option Yes selected, then go to question 4
- No other options lead to the requested service

4. Choose all that apply:⁽⁸⁾

- A) No septicemia
- B) No active osteomyelitis of the target vertebra
- C) No coagulopathy or coagulopathy treated
- D) No known allergy to bone cement or opacification agent
- E) Other clinical information (add comment)

30. Vertebral hemangioma (*Continued...*)

- If the number of options selected is 4 and option E not selected, then the rule is satisfied; you may stop here (*Outpatient*)
 - No other options lead to the requested service
-

Notes

(1)

I/O Setting: Outpatient

(2)

These criteria cover vertebroplasty and kyphoplasty in either the thoracic or lumbar spine.

(3)

Vertebroplasty is a percutaneous technique performed under fluoroscopic or CT guidance that involves the injection of polymethylmethacrylate cement into a collapsed vertebral body. It is performed for pain relief and to stabilize the fracture. Kyphoplasty, a modification of vertebroplasty, involves the insertion of an inflatable balloon within the vertebral body to expand the compressed vertebrae prior to cement injection. Kyphoplasty has been shown to restore vertebral body height and aid in the correction of kyphotic deformity that results from vertebral collapse (Kim et al., *J Spinal Disord Tech* 2012, 25: 338-44).

(4)

InterQual® Procedures criteria are derived from the systematic, continuous review and critical appraisal of the most current evidence-based literature and include input from our independent panel of clinical experts. To generate the most appropriate recommendations, a comprehensive literature review of the clinical evidence was conducted. Sources searched included PubMed, Agency for Healthcare Research and Quality (AHRQ) Comparative Effectiveness Reviews, the Cochrane Library, Choosing Wisely, Centers for Medicare & Medicaid Services (CMS) National Coverage Determinations, the National Institute of Health and Care Excellence (NICE), and the National Guideline Clearinghouse. Other medical literature databases, medical content providers, data sources, regulatory body websites, and specialty society resources may also have been used. Relevant studies were assessed for risk of bias following principles described in the Cochrane Handbook. The resulting evidence was assessed for consistency, directness, precision, effect size, and publication bias. Observational trials were also evaluated for the presence of a dose-response gradient and the likely effect of plausible confounders.

(5)

Spinal neurologic deficits are sensory or motor abnormalities due to neurocompression of either the spinal cord or nerve root and can include muscle weakness, paralysis, pain, or paresthesias. When present, decompression of the affected nerve should be considered.

(6)

Malignancy-related compression fractures are often difficult to distinguish from osteoporotic compression fractures, since clear radiographic findings of the malignancy are sometimes absent. These criteria are, therefore, satisfied when malignancy has been documented by biopsy, a history of recent malignancy, or radiographic evidence of a malignancy (Sutcliffe et al., *Health Technol Assess* 2013, 17: 1-274).

(7)

Since malignancy-related compression fractures are likely to progress, a trial of conservative treatment is not advised (Sutcliffe et al., *Health Technol Assess* 2013, 17: 1-274).

(8)

The listed findings are absolute contraindications to vertebroplasty or kyphoplasty (American College of Radiology (ACR), ACR Appropriateness Criteria: Management of Vertebral Compression Fractures. 2013).

(9)

The pain associated with vertebral compression fractures is usually intense and sometimes incapacitating for the first few weeks, improving gradually over the ensuing months. If vertebroplasty or kyphoplasty are recommended, they should be used as a second-line procedure for patients who have failed or cannot tolerate conservative management (Baerlocher et al., *J Vasc Interv Radiol* 2014, 25: 165-70; Barr et al., *J Vasc Interv Radiol* 2014, 25: 171-81; American College of Radiology (ACR), ACR Appropriateness Criteria: Management of Vertebral Compression Fractures. 2013).

(10)

NSAIDs are the preferred treatment of this condition because of their anti-inflammatory effect and are recommended in national clinical guidelines (Goertz et al., *Adult Acute and Subacute Low Back Pain*. 2012; Koes et al., *Eur Spine J* 2010, 19: 2075-94). Although evidence does not support the use of acetaminophen as first line therapy for treating spine pain, it can be used as an alternative when NSAIDs are contraindicated (e.g., pregnancy, history of peptic ulcer disease) (Machado et al., *BMJ* 2015, 350: h1225).

(11)

A back brace promotes pain relief from vertebral compression fractures by stabilizing the spine. Bracing limits pain-provoking movements and can facilitate early mobilization (Longo et al., *J Bone Joint Surg Br* 2012, 94: 152-7).

(12)

Activity modification involves limiting activities which provoke or aggravate symptoms. Patients may require bed rest in severe cases, although the risks of prolonged immobility must be balanced against the symptomatic benefits.

(13)

Def: A pathologic fracture is a fracture that occurs when a bone is weakened by disease (e.g., osteoporosis, benign or malignant tumors, osteomyelitis).

ICD-9 (circle all that apply): 170.2, 198.5, 228.09, 733.13, 81.65, Other_____

ICD-10-CM (circle all that apply): C41.2, C79.51, C79.52, D18.09, M48.50XA, M80.08XA, M84.48XA, M84.68XA, Other_____

ICD-10-PCS (circle all that apply): 0PU33JZ, 0PU34JZ, 0PU43JZ, 0PU44JZ, 0QU03JZ, 0QU04JZ, 0QU13JZ, 0QU14JZ,
Other_____

CPT® (circle all that apply): 22510, 22511, 22512, Other_____